


NCEP Contributions to the WMO Severe Weather Forecasting Demonstration Project (SWFDP) and to the African Monsoon Multidisciplinary Analysis (AMMA) Initiative

1.0. Rainfall Forecast: Valid 06Z of 18 January – 06Z of 22 January 2012, (Issued at 17:30Z of 17 January 2012)

1.1. Twenty Four Hour Cumulative Rainfall Forecasts

The forecasts are expressed in terms of 75% probability of precipitation (POP) exceeded, based on the NCEP, UK Met Office and the ECMWF NWP outputs, the NCEP global ensemble forecasts system (GEFS) and expert assessment.



<u>Summary</u>

In the next five days, seasonal wind convergence in the CAB region, localized wind convergences and cyclonic circulations in the vicinity of Angola, Namibia and Mozambique Channel are expected to enhance rainfall in their respective regions. Hence, there is an increased chance for heavy rainfall over eastern Angola, northern Namibia, Zambia, Zimbabwe, Botswana, southern DRC, Malawi, Mozambique, eastern South Africa and Madagascar.

1.2. Models Comparison and Discussion-Valid from 00Z of 17 January 2012

The GFS, ECMWF and UKMET models indicate series of lows and their associated troughs across central and the South African countries. The low over DRC is expected to deepen, with its mean sea level pressure value decreasing from 1008mb to 1005mb while shifting slightly to the south towards end of the forecast period according to the GFS model. According to ECMWF model, it is expected to deepen from MSLP value decreasing from 1007mb to 1006mb while shifting to the northwest towards end of the forecast period. According to the UKMET model, it is expected to deepen, with its MSLP value decreasing from 1008mb to 1007mb through 24 to 96 hours and then it tends to fill up back to 1008mb towards end of the forecast period. Another low is expected to form in the vicinity of Botswana and tends to fill up, with its MSLP value increasing from 1005mb to 1006mb through 24 to 72 hours and then it tends to deepen back to 1003mb while shifting slightly to the south towards end of the forecast period according GFS model. According to ECMWF model, the low pressure is expected to maintain its MSPL value of 1008mb towards end of the forecast period. This low pressure is expected to fill up, with its MSLP value increasing from 1005mb to 1008mb through 24 to 72 hours while shifting slightly to the south and then it tends to deepen, with its MSLP value decreasing from 1008mb to 1006mb and then it tends to fill up back to 1008mb towards end of the forecast period according UKMET model. Another low pressure is expected to form across Mozambique Channel and tends to deepen, with its mean sea level pressure value decreasing from 1003mb to 980mb while shifting to the southeast and reaching Madagascar towards end of the forecast period, according GFS model. According to the ECMWF, this low pressure is expected to deepen, with its MSLP value decreasing from 1008mb to 998mb while shifting slightly to the southeast towards end of the forecast period. This low tends to deepen, with its MSPL value decreasing from 1005mb to 995mb towards end of forecast period, according UKMET model. The fourth low over Sudan tends to fill up, with its MSLP value increasing from 1006mb to 1007mb through 24 to 72 hours and then it tends to deepen back to 1004mb towards end of the forecast period, according to GFS model. This low pressure is expected to fill up, with its MSLP value increasing from 1007 to 1008mb while shifting slightly to the south towards end of the forecast period according to ECMWF model. According the UKMET model, this low pressure tends to increase from 1004mb to 1006mb through 24 to 72 hours and then it tends to deepen back to 1005mb by 96 hours and then it tends to fill up back to 1008mb until the end of the forecast period.

The St. Helena High pressure system over southeast Atlantic Ocean is expected to fill up, with its MSLP value increasing from 1016mb to 1024mb through 24 to 96 hours and then it tends to deepen back to 1020mb while shifting slightly to the east and towards end of the forecast period according to **GFS** model. According to **ECMWF** model this high is expected to fill up, with its MSLP value increasing from 1016mb to 1024mb while moving to the east towards end of the forecast period. According to **UKMET** model, this high pressure tends to fill up, with its MSLP value increasing from 1160mb to 1028mb towards end of the forecast period. The Mascarene high pressure system over southwest Indian Ocean is expected to deepen with its central pressure value decreasing from 1020mb to 1016mb towards the end of the forecast period according to **GFS** model. According to **ECMWF** model it is expected to deepen, with its MSLP value decreasing from 1020mb to 1012mb towards end of the forecast period. The forecast period according to **GFS** model. According to **ECMWF** model it is expected to deepen, with its MSLP value decreasing from 1020mb to 1012mb towards end of the forecast period. This high pressure is expected to deepen, with MSLP value decreasing from 1020mb to 1012mb towards end of the forecast period. This high pressure is expected to deepen, with MSLP value decreasing from 1020mb to 1012mb towards end of the forecast period.

At the 850hpa level, a lower tropospheric seasonal wind convergence is expected to remain active over the CAB region aligning itself with a convergence across southern Africa. The convergence tends to intensify while shifting slightly to the southwest and reach Zambia, Malawi through 24 to 48 hours and then Zimbabwe and Mozambique towards end of the forecast period. Localized wind convergences are also expected to dominate the flow over Angola and Namibia then it tends to intensify and to shift slightly to the east towards end of the forecast period. Another localized wind convergences is expected to dominate the flow over Congo and Gabon then it tends to intensify and to shift slightly and to shift towards west towards end of forecast period. Lower tropospheric anticyclone and its associated ridge are expected over South Africa and the neighboring areas and tend to intensify and shift slightly to the west towards end of the forecast period. Another lover tropospheric cyclone and its associated trough are expected to dominate the flow over Mozambique towards end of the forecast period. The neighboring areas and tend to intensify and shift slightly to the west towards end of the forecast period. Another lover tropospheric cyclone and its associated trough are expected to dominate the flow over Mozambique Channel and it tends to shifting to the east and reach coastal Madagascar towards end of the forecast period.

At 500hpa, eastward propagating trough in the mid-latitude westerly flow is expected to prevail over Mediterranean Sea and northern Africa during the forecast period; with the low geopotential value of 5640gpm extending to the latitudes of Algeria by 24hours. This northeast-southwest oriented trough is expected to move southwards until reaching Mali and the adjoining areas with the low geopotential value of 5760gpm towards end of the forecast period. A mid latitude frontal system is also expected to propagate eastwards across the Southern African countries with the low geopotential value of 5820mb through 96 to 120 hours. Another trough is expected to prevail over Mozambique Channel by 48 hours with the low geopotential value of 5700gpm towards end of the forecast period.


At 200mb, strong winds associated with Sub-Tropical Westerly Jet are expected to dominate the flow over northern Africa, during the forecast period. The intensity of the jet is expected to exceed 150kts in the region between Atlantic Ocean and the Persian Gulf across Mauritania, southern Algeria, Libya and Egypt while moving to the east towards end of the forecast period. The Sub-tropical Westerly Jet in the Southern Hemisphere is expected to dominate the flow over the South Africa, by 120 hours. The intensity of the jet is expected to exceed 110kts in the region between southeastern Atlantic Ocean and South Africa.

In the next five days, seasonal wind convergence in the CAB region, localized wind convergences and cyclonic circulations in the vicinity of Angola, Namibia and Mozambique Channel are expected to enhance rainfall in their respective regions. Hence, there is an increased chance for heavy rainfall over eastern Angola, northern Namibia, Zambia, Zimbabwe, Botswana, southern DRC, Malawi, Mozambique, eastern South Africa and Madagascar.

2.0. Previous and Current Day Weather Discussion over Africa

(16 January – 17 January 2012)

- 2.1. Weather assessment for the previous day (16 January 2012): During the previous day, moderate to locally heavy rainfall was observed over northern Morocco, parts of Angola, Malawi and Mozambique.
- **2.2. Weather assessment for the current day (17 January 2012):** Intense clouds are observed over DRC, Zambia, eastern South Africa, Angola, Namibia, Malawi, Tanzania, northern Mozambique and Madagascar.



Previous day rainfall condition over Africa (top Left) based on the NCEP CPCE/RFE and current day cloud cover (top right) based on IR Satellite image

Author(s): Lotfi Khammari (Tunsian Meteorological Authority) / CPC-African Desk), lotfi.khammari@noaa.gov, and

Author(s): Amira Ibrahim (Egyptian Meteorological Authority) / CPC-African Desk), <u>Amira.ibrahim@noaa.gov</u>