

1 **Title: Current Atlantic Meridional Overturning Circulation weakest in last**  
2 **millennium**

3 **Authors:** L. Caesar<sup>1,2\*</sup>, G. D. McCarthy<sup>1</sup>, D. J. R. Thornalley<sup>3</sup>, N. Cahill<sup>4</sup>, S. Rahmstorf<sup>2,5</sup>

4 **Affiliations:**

5 <sup>1</sup>Irish Climate Analysis and Research Units (ICARUS), Department of Geography, Maynooth  
6 University, Maynooth, Ireland.

7 <sup>2</sup>Potsdam Institute for Climate Impact Research (PIK), Member of the Leibniz Association, P.O.  
8 Box 60 12 03, D-14412 Potsdam, Germany.

9 <sup>3</sup>Department of Geography, University College London, London, UK.

10 <sup>4</sup>Department of Mathematics and Statistics, Maynooth University, Kildare, Ireland.

11 <sup>5</sup>Institute of Physics and Astronomy, University of Potsdam, Potsdam, Germany.

12 **Abstract – The Atlantic meridional overturning circulation (AMOC)— one of Earth’s**  
13 **major ocean circulation systems— redistributes heat on our planet and has a major impact**  
14 **on climate. Here, we compare a variety of published proxy records to reconstruct the**  
15 **evolution of the AMOC since about 400 AD. Taken together these data presents a fairly**  
16 **consistent picture of the AMOC: After a long and relatively stable period follows an initial**  
17 **decline in the AMOC starting in the 19th Century, with a second, more rapid, decline**  
18 **following in the mid-20th Century. Taken together, these data suggest that, during the last**  
19 **decades, the AMOC has been at its weakest state for over a millennium.**

20 The Atlantic Meridional Overturning Circulation (AMOC) is a major mechanism for heat  
21 redistribution on our planet and an important factor in climate variability and change. The  
22 AMOC is a sensitive non-linear system dependent on subtle thermohaline density differences in  
23 the ocean, and major AMOC transitions have been implicated e.g. in millennial climate events  
24 during the last glacial period <sup>1</sup>. There is evidence that the AMOC is slowing down in response to  
25 anthropogenic global warming <sup>2</sup>—as predicted by climate models—and that the AMOC is  
26 presently in its weakest state for more than 1000 years <sup>3</sup>. As continuous direct measurements of  
27 the AMOC only started in 2004 <sup>4</sup>, longer term reconstruction must be based on proxy data. In  
28 general, there are three different types of AMOC proxies: i) reconstructions of surface or  
29 subsurface temperature patterns in the Atlantic Ocean that reflect the changes in ocean heat  
30 transport associated with the AMOC <sup>3,5</sup>; ii) reconstructions of subsurface water mass properties,  
31 e.g. the advance of the subpolar vs subtropical slope water, that reflect AMOC changes e.g. <sup>6</sup>;  
32 and iii) evidence for physical changes in deep-sea currents, such as those reflected by changes in  
33 sediment grain size <sup>3</sup>. As all kinds of proxies are limited in their representation of the AMOC (all  
34 three can be influenced to some degree by factors in addition to changes in the AMOC), a  
35 combination of all three proxy types is needed to provide robust evidence about the evolution of  
36 the AMOC.

37 Here, using several different and largely independent proxy indicators of the AMOC evolution  
38 over the last one hundred to nearly two thousand years, we provide strong evidence that the  
39 AMOC decline in the 20th Century is unprecedented and that over the last decades the AMOC is  
40 in its weakest state in over a millennium.

41 The proxies are taken from various locations in the Atlantic or the surrounding land areas (inset  
42 of figure 1) and represent either different subsystems associated with the AMOC (like Labrador

43 Sea density<sup>3</sup>, the presence of subtropical versus subpolar slope waters along the North American  
44 East coast<sup>6,7</sup>) or the effect of changes in the Atlantic meridional heat transport associated with  
45 the AMOC<sup>2,3,5,8</sup>, as well as surface ocean productivity changes that have been related to the  
46 AMOC<sup>9,10</sup>. The records going the furthest back in time (400 AD) are taken from marine  
47 sediments (sortable-silt data<sup>3</sup>, proxy records of subsurface ocean temperatures<sup>3</sup>,  $\delta^{18}\text{O}$  in benthic  
48 foraminifera<sup>7</sup>,  $\delta^{15}\text{N}$  of deep-sea gorgonian corals<sup>6</sup>, relative abundance of certain planktic  
49 foraminifera (*Turborotalita quinqueloba*)<sup>10</sup>). The temperature-based AMOC index<sup>5</sup> on the other  
50 hand is based on a Northern Hemisphere land-and-ocean temperature reconstruction that uses a  
51 range of terrestrial proxies including e.g. tree rings and ice core data<sup>11</sup>. Data taken from  
52 Greenland ice cores (the methanesulfonic acid concentration) furthermore provide an estimate  
53 for AMOC related changes in productivity in the subpolar gyre (SPG) region<sup>9</sup>. Most of these  
54 records extend into the modern era, for which additional AMOC proxies exist that are based on  
55 instrumental temperature records<sup>2,8</sup>.

56 Despite the different locations, time scales and processes represented by these proxies, they  
57 provide a consistent picture of the AMOC evolution since about 400 AD: Prior to the 19<sup>th</sup>  
58 century, the AMOC was relatively stable. A decline in the AMOC, beginning during the 19<sup>th</sup>  
59 century, is evident in all the proxy records (figure 1 left panel). Around 1960 a phase of  
60 particularly rapid decline started that is found in several, largely independent proxies. A short-  
61 lived recovery is evident in the 1990s before a return to decline from the mid-2000s (figure 1  
62 right panel). All indices additionally show multi-decadal variability, albeit with different  
63 amplitudes and frequencies making it questionable whether this is mainly driven by the AMOC.  
64 Some of the differences likely relate to the large range in temporal resolution in the proxies  
65 (from annual to 50-year binning), while others are likely due to complicating factors, such as

66 non-AMOC related influences on a proxy system (e.g. changes in trophic structure of coral's  
67 food source in  $\delta^{15}\text{N}$ , local fluctuations in circulation impacting single site palaeoceanographic  
68 reconstructions, or other controls on subpolar heat content <sup>12</sup>). An additional factor may also be  
69 that different components of the AMOC respond on different time scales. While the strength of  
70 the AMOC, typically measured at 26°N, has been shown to be correlated to the multi-decadal  
71 variability of North Atlantic SST <sup>13</sup> (suggesting that a large part of this variability in the  
72 temperature-based proxies are due to AMOC changes) changes in the deep ocean appear to occur  
73 on a different timescale. Therefore, it is unsurprising that for the larger part of the last  
74 millennium the multi-decadal variability in the proxies differ.

75 The strength of this multi-proxy comparison lies in tracing the centennial and longer AMOC  
76 evolution. To test whether the reduction in AMOC strength that is seen in all proxy records is  
77 significant, a change-point model is fitted to each time series and the data means before and after  
78 the change point are compared (see Methods). Assuming, in the first approximation, only a  
79 single change point, the model finds a significant reduction in the mean in all but one proxy  
80 record (see table 1). The timing of the change point varies in the different proxy series (also due  
81 to the different lengths of the time series) but can be sorted into two clusters: one change  
82 occurring in the second half of the 19<sup>th</sup> century and a second change occurring in the 1960s. To  
83 test the significance of differences between different time periods, we divided each time series  
84 into 50 year intervals (30 year intervals for the Cheng et al. (2017) data given that the length of  
85 the time series is only 64 years and 100 year intervals for the Spooner et al. (2020) data given the  
86 coarse resolution of this time series), going backward from the present and we estimated the  
87 means and data uncertainty for each of these intervals. The mean of any 50 (30, 100) year  
88 interval is assumed to be significantly lower when its uncertainty range does not overlap with the

89 uncertainty range of the mean of any other interval. The results show that in 9 of the 11 proxy  
90 series the most recent 50 (30, 100) year mean value is significantly lower than any other before  
91 (see table 1). In addition, the high-resolution proxies suggest a progressive AMOC decline  
92 within that most recent interval.

93 Together these data consistently show that the modern AMOC slowdown is unprecedented in  
94 over a thousand years. Improved understanding of this slowdown is urgently needed. The next  
95 step is to resolve which components and pathways of the AMOC have altered, how, and why -  
96 no small feat, and requiring a community effort that combines observational, modelling and  
97 palaeoclimatological approaches.

## 98 **Methods**

99 Uncertainties. The uncertainty range represents in all but one case the 2- $\sigma$  confidence interval for  
100 the individual proxy reconstruction, i.e. for i) the proxy-based surface temperature reconstruction  
101 (validated against independent instrumental temperature data)<sup>5</sup>, ii) the subsurface temperature  
102 dipole in the Atlantic based on the published uncertainties for age assignment and temperature  
103 reconstructions<sup>3</sup>, iii) the  $\delta^{15}\text{N}$  record based on a mixed effect linear model based on year and  
104 specimen colony<sup>6</sup>, iv) the sortable silt data that is shown with its full (reduced) procedural error  
105<sup>3</sup>, v) the  $\delta^{18}\text{O}$  data based on analytical reproducibility determined by replicate measurements of  
106 internal standard carbonate material<sup>7</sup>, vi) the abundance of *T. quinqueloba* with the uncertainty  
107 estimated using a binomial approach<sup>10</sup>, and vii) the marine productivity in the subpolar gyre  
108 based on a bootstrapping method<sup>9</sup>. As an upper bound for the 2- $\sigma$  confidence interval of the  
109 relative change in Atlantic Ocean heat content vs that in the Southern Ocean the confidence  
110 intervals for the individual ocean heat content time series, considering among others instrumental  
111 errors, methodological choices and data gaps, were simply added<sup>8</sup>.

112 For the temperature-based AMOC proxy <sup>2</sup> the uncertainty in converting this proxy data to an  
113 AMOC slowdown is given, not the uncertainty of the temperature data itself. This is based on the  
114 relationship between the relative temperature change in the subpolar North Atlantic and AMOC  
115 variability in the CMIP5 model ensemble.

116 Although only this last uncertainty interval considers the spread in the proxy that is unrelated to  
117 the AMOC, the other proxies have all been related to AMOC variability. Moreover, given that  
118 the proxies were taken from multiple locations across the Northern Hemisphere and the only  
119 inferred common driver for them all is AMOC, combined, they provide strong evidence for a  
120 centennial decline related to the AMOC.

121 Statistical significance. To determine whether there has been a significant change in the proxy  
122 time series that would indicate a reduction in AMOC strength we tested each proxy record for a  
123 single significant change in the mean of the time series. Using a Bayesian framework, we fit a  
124 model that assumes that the data fluctuate around a constant mean, allowing for a single change  
125 in the mean at some point in time. The approach takes both the data uncertainty and the data  
126 variability into account. Once the model finds the timing of the change, we compare the means  
127 before and after the change point to check if the difference is significant (we assume significance  
128 when the 95% Bayesian credible interval of the difference between the means does not contain a  
129 zero value).

130 To test whether the AMOC is at its weakest in over a millennium we applied a similar  
131 framework that fixed change points at 50 (30, 100) year intervals. Starting in the present, the  
132 mean and 95% uncertainty interval for each 50 (30, 100) year interval was estimated (taking data  
133 uncertainty and the number of data points in each interval into account).

134 **Data Availability**

135 The datasets analysed during the current study were provided by the authors from the original  
136 publications (see labels of figure 1). They are available from the corresponding author on  
137 request.

138 **Code Availability**

139 The script for analysing and plotting the data is available from the corresponding author upon  
140 request.

141 **Competing Interests statement:**

142 The authors declare no competing interests.

143 **References:**

- 144 1 Rahmstorf, S. Ocean circulation and climate during the past 120,000 years. *Nature* **419**,  
145 207-214, doi:10.1038/nature01090 (2002).
- 146 2 Caesar, L., Rahmstorf, S., Robinson, A., Feulner, G. & Saba, V. Observed fingerprint of  
147 a weakening Atlantic Ocean overturning circulation. *Nature* **556**, 191-196,  
148 doi:10.1038/s41586-018-0006-5 (2018).
- 149 3 Thornalley, D. J. R. *et al.* Anomalously weak Labrador Sea convection and Atlantic  
150 overturning during the past 150 years. *Nature* **556**, 227-230, doi:10.1038/s41586-018-  
151 0007-4 (2018).
- 152 4 Smeed, D. A. *et al.* The North Atlantic Ocean Is in a State of Reduced Overturning.  
153 *Geophysical Research Letters* **45**, 1527-1533, doi:10.1002/2017gl076350 (2018).
- 154 5 Rahmstorf, S. *et al.* Exceptional twentieth-century slowdown in Atlantic Ocean  
155 overturning circulation. *Nature Climate Change* **5**, 475-480, doi:10.1038/nclimate2554  
156 (2015).
- 157 6 Sherwood, O. A., Lehmann, M. F., Schubert, C. J., Scott, D. B. & McCarthy, M. D.  
158 Nutrient regime shift in the western North Atlantic indicated by compound-specific  
159 delta15N of deep-sea gorgonian corals. *Proc Natl Acad Sci U S A* **108**, 1011-1015,  
160 doi:10.1073/pnas.1004904108 (2011).

- 161 7 Thibodeau, B. *et al.* Last Century Warming Over the Canadian Atlantic Shelves Linked  
162 to Weak Atlantic Meridional Overturning Circulation. *Geophysical Research Letters* **45**,  
163 12,376-312,385, doi:10.1029/2018gl080083 (2018).
- 164 8 Cheng, L. *et al.* Improved estimates of ocean heat content from 1960 to 2015. *Science*  
165 *Advances* **3**, e1601545, doi:10.1126/sciadv.1601545 (2017).
- 166 9 Osman, M. B. *et al.* Industrial-era decline in subarctic Atlantic productivity. *Nature* **569**,  
167 551-+, doi:10.1038/s41586-019-1181-8 (2019).
- 168 10 Spooner, P. T. *et al.* Exceptional 20th Century Ocean Circulation in the Northeast  
169 Atlantic. *Geophysical Research Letters* **47**, e2020GL087577,  
170 doi:10.1029/2020GL087577 (2020).
- 171 11 Mann, M. E. *et al.* Proxy-based reconstructions of hemispheric and global surface  
172 temperature variations over the past two millennia. *Proceedings of the National Academy*  
173 *of Sciences of the United States of America* **105**, 13252-13257,  
174 doi:10.1073/pnas.0805721105 (2008).
- 175 12 Keil, P. *et al.* Multiple drivers of the North Atlantic warming hole. *Nature Climate*  
176 *Change* **10**, 667-671, doi:10.1038/s41558-020-0819-8 (2020).
- 177 13 Zhang, R. *et al.* A Review of the Role of the Atlantic Meridional Overturning Circulation  
178 in Atlantic Multidecadal Variability and Associated Climate Impacts. *Reviews of*  
179 *Geophysics* **57**, 316-375, doi:10.1029/2019RG000644 (2019).

180

181 **Corresponding author**

182 Correspondence to: Levke Caesar, levke.caesar@mu.ie

183 **Acknowledgments**

184 L. Caesar, N. Cahill and G. D. McCarthy are supported by the A4 project. A4 (Grant-Aid  
185 Agreement No. PBA/CC/18/01) is carried out with the support of the Marine Institute under the  
186 Marine Research Programme funded by the Irish Government, co-financed by the European  
187 Regional Development Fund. D. J. R. Thornalley is supported by UK NERC grant  
188 NE/S009736/1.



189 **Author contributions**

190 S.R. initiated the study. L.C. created the figure and wrote the manuscript. N.C. performed the  
191 significance testing. All authors discussed and interpreted the results and provided input to the  
192 manuscript.

193 **Figure Legends**

194 **Fig. 1: SST-based AMOC reconstructions compared to various proxy reconstructions. a,**

195 The SST-based proxies (light and dark blue) represent the North Atlantic temperature response  
196 to changes in the Atlantic meridional heat transport associated with an AMOC slowdown. It is  
197 compared to proxy records of **b**, subsurface ocean temperatures (purple), **c** and **h**,  $\delta^{15}\text{N}$  data of  
198 deep-sea gorgonian corals (magenta), **d** and **i**, sortable-silt data (shades of green, shown with a  
199 12-year lag to the temperature-based indices <sup>3</sup>), **e** and **j**,  $\delta^{18}\text{O}$  data in benthic foraminifera (shades  
200 of brown), **f**, the relative abundance of *T. quinqueloba* in marine sediment cores (orange-red) as  
201 well as, **f** and **k**, methanesulfonic acid concentration in Greenland ice cores (orange), both  
202 indicators for local/regional marine productivity, and **g**, the relative change in Atlantic Ocean  
203 heat content vs that in the Southern Ocean (dark magenta, only in the right panel). As a reference  
204 for the actual change in volume transport the April 2004 – April 2018 linear trend of the RAPID  
205 data <sup>4</sup> (black) is given (**g**). The map (using the same color-coding as the time series) gives an  
206 overview of the various locations the proxies were taken from (with small markers denoting  
207 single sites and large markers denoting the areas with multiple proxy sites). All curves were  
208 smoothed with a 20-year (50-year) LOWESS filter for the shorter (longer) time series to make  
209 them more comparable. Shading and error bars show the  $2\sigma$ -(95%)-confidence interval of the

210 individual proxies as they were reported and the uncertainty of the AMOC representation of the  
 211 Caesar et al. (2018) temperature proxy, respectively (see Methods).

212 **Tables**

General Information			Change point testing		Significance testing	
Proxy	Time interval	Long/Short	95%-interval	Signif. reduction	Lowest interval	Signif. lower
Temperature anomaly <sup>5</sup>	900-1995	L	1874-1902	yes	1946-1995	yes
Subsurface temperature proxy <sup>3</sup>	400-2000	L	1817-1856	yes	1951-2000	yes
$\delta^{15}\text{N}$ data <sup>6</sup>	1926-2002*	S	1970-1976	yes	1953-2002	yes
Sortable silt data 48JPC <sup>3</sup>	380-1995 <sup>†</sup>	L	1763-1878	yes	1876-1925	no
Sortable silt data 56 JPC <sup>3</sup>	1475-2003	L	1863-1883	yes	1904-1953	no
$\delta^{18}\text{O}$ data <sup>7</sup>	708-1962	L	1881-1916	yes	1913-1962	yes
<i>T. quinqueloba</i> abundance <sup>10</sup>	392-2013	L	1920-1958	yes	1914-2013	yes
Temperature proxy <sup>2</sup>	1871-2016	S	1967-1970	yes	1967-2016	yes
$\delta^{18}\text{O}$ data <sup>7</sup>	1904-2001	S	1960-1975	yes	1952-2001	yes
Marine productivity <sup>9</sup>	1767-2013	S	1950-1956	yes	1964-2013	yes
Ocean heat content <sup>8</sup>	1955-2019	S	For this data set the algorithm did not find a significant change point.		1990-2019	yes

\*  $\delta^{15}\text{N}$  data starts in 565 AD but is continues only from 1926 onwards.  
<sup>†</sup> The last data point of the 48JPC sortable silt data is in 1995, but due to robustness for the significance testing the smoothed data was used which extends only until 1975 (as this is the penultimate data point).

213 **Table 1: Results of the change point and significance testing of the various proxies used to**  
 214 **reconstruct the evolution of the AMOC.** The first three columns include general information  
 215 about the proxies like the covered time interval and the categorization into long (L) or short (S)  
 216 proxy time series. The columns in the middle list the 95%-interval of the change point found by

217 the change point model with most long time series and most short time series having a change  
218 point in the late 19<sup>th</sup> Century and in the 1960s, respectively. It is additionally noted whether the  
219 reduction in the proxy following the change point is significant. The columns at the right list the  
220 50 (30, 100) year interval during which the proxy is at its lowest value and whether this value is  
221 significantly low compared to all other 50 (30, 100) year intervals (considering data uncertainty).